

Standing Nurse Orders

East Kingston Elementary
Kensington Elementary
Newfields Elementary

Exeter High School
Lincoln Street School
Stratham Memorial

Cooperative Middle School
Main Street School
Swasey Central School

Opiate Overdose - grades 6-12

For suspected opioid overdose with unresponsiveness and poor or no respirations, Call 911.

May administer Naloxone (Narcan) 2mg/2ml: 1 ml spray in each nostril. May repeat if no response in 3-5 minutes

Localized/NonSystemic Allergic Reaction: May give:

Benadryl (diphenhydramine HCL) PO	Up to 40 lb. -- ---- 12.5 mg.
	40 lb. to 100 lb. ---- 25 mg
	Over 100 lb. ----- 50 mg

Systemic Allergic Reaction: Difficulty breathing, hive like rash, facial or extensive body swelling

1. Benadryl (diphenhydramine HCL) PO given if able to swallow - see dosage above
2. Give EPI-PEN (Adrenaline chloride 1/1000) IM by estimated body weight:
Up to 60 lb. - (age 8) Epi-Pen Jr. (0.15mg)
Over 60 lb. - (over age 8) – Epi-Pen Adult (0.3mg)
3. This may be repeated after 20 minutes if necessary
4. Call 911 and parent after an Epi-Pen is given

Oxygen:

May administer oxygen via nasal cannula at 2-6 liters if oxygen is available and appropriate to the situation.

Severe Hypoglycemia: unable to take food or drink by mouth

Glucagon: 1 mg IM (1/2 mg IM if student weighs <44 pounds)

Glucagon may cause nausea and vomiting; place student on his side.

Notify parents after giving glucagons.

Seasonal Allergic Rhinitis: Grades 6-12 only

Loratadine 10 mg po with parent permission:

Minor Aches, Discomfort, Headaches, Fever, Menstrual Cramps:

Advil (Ibuprofen) or Tylenol (Acetaminophen) as determined by age/weight every four to six hours PRN with Parent Permission

The following items (or generic equivalent) may be used for the purpose of rendering first aid and comfort based on standard nursing practice **unless otherwise indicated by parent**: cold or heat application, bacitracin, triple antibiotic, hydrocortisone cream, caladryl, sting kill swabs, burn gel, anbesol, chloraseptic spray, throat lozengers, cough drops, basic antacid

Per order of

W. J. Lee mo 9/19/16
Doctor
Michelle Moya 9/20/16
Superintendent date

Standing Nurse Orders Child Asthma

East Kingston Elementary
Exeter High School
Cooperative Middle School

Kensington Elementary
Lincoln Street School
Main Street School

Newfields Elementary
Stratham Memorial
Swasey Central School

Asthma:

In the case of a child with asthma - wheezing and shortness of breath - and if the child does not have their own inhaler with them at school, the nurse should:

1. Assess the status of air exchange of the child via

Vital signs

Listen to the chest with a stethoscope

Peak flow meter

2. If the peak flow is above 80% of the child's predicted peak flow, try to relieve the child's anxiety and observe
3. If the peak flow is 50% to 80% of the child's predicted peak flow, administer Albuterol inhaler

MDI Albuterol - 2 puffs separated by two minutes

4. If the peak flow is below 50% of the child's predicted peak flow, administer a nebulizer and refer to the ER if unable to reach parents and the child does not immediately respond to the nebulizer.

Nebulizer with Albuterol 0.083% with NS diluent.

Per order of _____
Doctor *[Signature]* date 9/19/16
Superintendent *[Signature]* Date 9/20/16